

Analyses of multiplicity distributions of e^+e^- and $e-p$ collisions by means of modified negative binomial distribution and Laguerre-type distribution: Interrelation of solutions in stochastic processes

Minoru Biyajima¹, Takeshi Osada² and Kenji Takei¹

¹Department of Physics, Shinshu University, Matsumoto 390, Japan

²Department of Physics, Tohoku University, Sendai 980-77, Japan

February 1, 2008

Abstract

A pure birth stochastic process with several initial conditions is considered. We analyze multiplicity distributions of e^+e^- collisions and $e-p$ collisions, using the Modified Negative Binomial Distribution (MNBD) and the Laguerre-type distribution. Several multiplicity distributions show the same minimum χ^2 's values in analyses by means of two formulas: In these cases, we find that a parameter N contained in the MNBD becomes to be large. Taking large N limit in the MNBD, we find that the Laguerre-type distribution can be derived from it. Moreover, from the generalized MNBD we can also derive the generalized Glauber-Lachs formula.

Finally stochastic properties of QCD and multiparticle dynamics are discussed.

PACS number(s): 13.65.+1, 34.10.+x

1. Introduction

The Negative Binomial Distribution (NBD) played the important role in analyses of multiplicity distributions in high energy collisions[1, 2]. However, to resolve the problem concerning a parameter k of the NBD the following scheme adopted from QCD was proposed in Refs. [3, 4]. In QCD we can consider a pure gluon bremsstrahlung process, $g \rightarrow g + g$. The evolution equation is

$$\frac{\partial P(n, t)}{\partial t} = -\lambda n P(n, t) + \lambda(n-1)P(n-1, t), \quad (1)$$

where $P(n, t)$, $t \propto \ln[\ln(Q^2/\mu^2)/\ln(Q_0^2/\mu^2)]$, and $\lambda = N_c(\text{color factor})/\varepsilon(\text{cutoff})$ are the gluon distribution, a proper variable in QCD, and the production rate of gluons, respectively. Assuming n_0 gluons at $t = 0$, i.e., δ_{n,n_0} , they obtained the following distribution with $\bar{n}(t) = n_0 \exp(\lambda t)$,

$$P_{n_0}(n, t) = \frac{(n-1)!}{(n-n_0)!(n_0-1)!} \left(\frac{n_0}{\bar{n}}\right)^{n_0} \left(1 - \frac{n_0}{\bar{n}}\right)^{n-n_0}. \quad (2)$$

Using the local parton duality between QCD and multiparticle dynamics Eq.(2) was applied to the data of high energy collisions in Refs. [3, 4]. However, Eq.(2) does not show satisfactory results, because of the fixed multiplicity at $t = 0$. This suggests that the fluctuation at $t = 0$ is important.

Later to improve the defects of Eq. (2), two initial conditions, the Poissonian distribution and the binomial distribution have been considered: Using Eq. (1) with them, the Laguerre-type distribution in Ref. [5] and the Modified Negative Binomial Distribution (MNBD) in Refs. [6]-[10] have been derived. See also Refs.[11]-[13].

In the present paper we analyze the data of e^+e^- collisions including the data of e - p collisions in Refs.[14]-[21] by means of three formulas. Several results show that the minimum χ^2 values by the MNBD with large N contained in the formula are coincided with those by the Laguerre-type distribution. See Table I. Thus we investigate whether or not there is any interrelation between the MNBD and the Laguerre-type distribution.

Contents of this paper are as follows: In the next paragraph and the third one, we introduce the MNBD and the Laguerre-type distribution, respectively. In the fourth one, we consider the large N limit of the MNBD. In the fifth one, we consider the generalized MNBD. Finally our concluding remarks are presented.

2. MNBD and its application to data of e^+e^- and e - p collisions

We assume that the solution of Eq.(1), i.e., $P(n, t)$, is useful for description of multiplicity, i.e., in multiparticle dynamics, using the local parton duality. The MNBD is obtained from Eq.(1), using the following initial condition,

$$P(n, t = 0) = {}_N C_n \alpha^n (1 - \alpha)^{N-n}, \quad (3)$$

where N is an integer and $N\alpha$ is the average number of clusters. The generating function of the MNBD is

$$\Pi(u) = \sum_{n=0}^{\infty} P(n) u^n = [1 - r_1(u-1)]^N [1 - r_2(u-1)]^{-N}, \quad (4)$$

where $r_1 = p - \alpha(p+1)$, and $r_2 = p$ with $p = \exp(\lambda t) - 1$. In this case λ is the production rate of hadrons.

The MNBD is given by the following function [6, 8], neglecting the second variable,

$$\begin{aligned}
P(0) &= \left[\frac{1+r_1}{1+r_2} \right]^N, \\
P(n) &= \frac{1}{n!} \left[\frac{1+r_1}{1+r_2} \right]^N \sum_{s=0}^n {}_n C_s \left[(-A)^s \frac{N!}{(N-s)!} \right] \\
&\quad \times \left[B^{n-s} \frac{(N+n-s-1)!}{(N-1)!} \right],
\end{aligned} \tag{5}$$

where $A = 1 + r_1/(1 + r_1)$ and $B = r_2/(1 + r_2)$. The r_1 and r_2 are expressed by the average multiplicity $\langle n \rangle$ and the second moment $C_2 = \langle n^2 \rangle / \langle n \rangle^2$. In this case r_1 is real and $r_2 > 0$

$$r_1 = \frac{1}{2} \left(C_2 - 1 - \frac{1}{N} \right) \langle n \rangle - \frac{1}{2}, \quad r_2 = r_1 + \frac{\langle n \rangle}{N}. \tag{6}$$

Our analyses of the charged multiplicity distributions in e^+e^- and e - p collisions are shown in Table I.

The KNO scaling function of the MNBD has been calculated by the usual procedure, $z = \lim_{n, \langle n \rangle \rightarrow \infty} n / \langle n \rangle$ fixed, in a recent paper [10]:

$$\Psi(z) \equiv \lim_{n, \langle n \rangle \rightarrow \infty} \langle n \rangle P(n) = \left(\frac{r'_1}{r'_2} \right)^N e^{-\frac{\langle n \rangle}{r'_2} z} \sum_{j=1}^N {}_N C_j \frac{1}{\Gamma(j)} \left(\frac{r'_2 - r'_1}{r'_1} \right)^j \left(\frac{\langle n \rangle}{r'_2} \right)^j z^{j-1}. \tag{7}$$

The parameters r'_1 and r'_2 in Eq. (7) are given by

$$r'_1 = \frac{1}{2} \left(C_2 - 1 - \frac{1}{N} \right) \langle n \rangle, \quad r'_2 = r'_1 + \frac{\langle n \rangle}{N}, \tag{8}$$

because $\langle n \rangle \gg 1$. Our results by the KNO scaling functions are given in Table II.

3. Laguerre-type distribution and its application to data

Using Eq. (1) with the Poissonian distribution as the initial condition,

$$P(n, t=0) = \delta_{n,k} \frac{\langle m \rangle^k}{k!} \exp(-\langle m \rangle), \tag{9}$$

we obtain the Laguerre-type distribution. (Notice that $\langle m \rangle = 2/(C_2 - 1 + 1/\langle n \rangle)$ has been interpreted as the number of clusters at $t=0$.)

$$\begin{aligned}
P(0) &= \exp(-\langle m \rangle), \\
P(n) &= \langle m \rangle^2 \exp(-\langle m \rangle) \frac{(\langle n \rangle - \langle m \rangle)^{n-1}}{\langle n \rangle^n} \frac{1}{n} L_{(n-1)}^{(1)} \left(-\frac{\langle m \rangle^2}{\langle n \rangle - \langle m \rangle} \right), \quad n = 1, 2, \dots
\end{aligned} \tag{10}$$

It should be stressed that Eq.(10) was derived in various approaches and applied to analyses of data in high energies in Refs. [11, 12, 13].

The KNO scaling function of Eq.(10) is given as

$$\Psi(z) = \frac{\langle m \rangle_k}{\sqrt{z}} \exp[-\langle m \rangle_k(z+1)] \times I_1(2\langle m \rangle_k \sqrt{z}), \quad (11)$$

where $\langle m \rangle_k = 2/(C_2 - 1)$. Our results by means Eqs. (10) and (11) are also given in Table I and II. Therein we can see that several minimum χ^2 's values by Eq.(5) with large N are coincided with those by Eq.(10). This fact suggests us that the MNBD in large N limit probably contains the Laguerre-type distribution. Thus in the next paragraph we investigate this possibility.

4. Large N limit of the MNBD

We take the large N limit in Eq. (5):

$$\left(\frac{1+r_1}{1+r_2}\right)^N \xrightarrow{\text{large } N} \exp\left(-\frac{2}{C_2 - 1 + 1/\langle n \rangle}\right) = \exp(-\langle m \rangle) \quad (12)$$

Eq. (12) is interpreted as the distribution of clusters at the initial time $t = 0$ as Ref. [5]. The origin is attributed to the Poissonian distribution at $t = 0$, which can be reduced from the binomial distribution as

$${}_N C_n \alpha^n (1-\alpha)^{N-n} \xrightarrow{\text{large } N} \frac{(N\alpha)^n}{n!} \exp(-N\alpha). \quad (13)$$

Using $N\alpha = \langle n \rangle / (1+r_2) = \langle m \rangle$, we obtain Eq.(10) as Ref. [5]. Using an approximation ${}_N C_j \approx N^j / j!$, we obtain Eq.(10) neglecting the contribution with $\mathcal{O}(1/N)$. Thus it can be said that Eq.(5) contains Eq.(10). For the KNO scaling functions, we have Eq.(11) taking the large N limit in Eq.(17), and using the following approximation

$$\lim_{N \rightarrow \infty} \frac{1}{N} L_{N-1}^{(1)}\left(\frac{-1}{N} \langle m \rangle_k^2 z\right) \approx \frac{1}{\langle m \rangle_k \sqrt{z}} I_1(2\langle m \rangle_k \sqrt{z}). \quad (14)$$

The present calculation shows that there is an interesting interrelation between solutions of Eq (1). Here it is worthwhile to notice that a similar interrelation between the Ising model with the free lattice and Eq.(10) was used in Ref.[12].

5. Generalized MNBD

According to calculations of Refs. [2,3], we can consider Eq.(1) with the immigration as follows

$$\partial P(n, t) / \partial t = -\lambda n P(n, t) + \lambda(n-1)P(n-1, t) - \lambda_0 P(n, t) + \lambda_0 P(n-1, t), \quad (15)$$

where λ_0 is an immigration rate corresponding to $q \rightarrow q + g$ process in QCD[2-3]. (On the other hand, in multiparticle dynamics the physical meaning of λ_0 may be interpreted as possible contributions from neutral excited hadrons.)

By making use the same initial condition, Eq. (3), we obtain the following generating function with $k = \lambda_0/\lambda$ as

$$\Pi(u) = \sum_{n=0} P(n)u^n = [1 - \tilde{r}_1(u-1)]^N [1 - \tilde{r}_2(u-1)]^{-k-N}, \quad (16)$$

where

$$\tilde{r}_1 = \frac{\langle n \rangle}{k} \left\{ 1 - \sqrt{\frac{k+N}{N} \left[1 - k \left(C_2 - 1 - \frac{1}{\langle n \rangle} \right) \right]} \right\}, \quad \tilde{r}_2 = \frac{N\tilde{r}_1 + \langle n \rangle}{k+N}.$$

Notice that Eq. (16) contains the NBD($N=0$) and the MNBD($k=0$). The probability distribution is obtained from $\Pi(u)$ as

$$\begin{aligned} P(0) &= \Pi(0) = \frac{(1 + \tilde{r}_1)^N}{(1 + \tilde{r}_2)^{N+k}}, \\ P(n) &= \frac{1}{n!} \frac{(1 + \tilde{r}_1)^N}{(1 + \tilde{r}_2)^{N+k}} \sum_{s=0}^n {}_n C_s \left[(-\tilde{A})^s \frac{N!}{(N-s)!} \right] \\ &\quad \times \left[\tilde{B}^{n-s} \frac{(N+k+n-s-1)!}{(N+k-1)!} \right] \end{aligned} \quad (17)$$

where $\tilde{A} = \tilde{r}_1/(1 + \tilde{r}_1)$ and $\tilde{B} = \tilde{r}_2/(1 + \tilde{r}_2)$.

We obtain the following generalized Glauber-Lachs formula in Ref.[1] taking large N limit in Eq.(17),

$$\begin{aligned} \text{Eq. (17)} \xrightarrow{\text{large } N} P(n) &= \frac{(p\langle n \rangle/k)^n}{(1 + p\langle n \rangle/k)^{n+k}} \exp\left[-\frac{\langle n \rangle}{(1 + p\langle n \rangle/k)}\right] \\ &\quad \times L_n^{(k-1)}\left(-\frac{k/p}{(1 + p\langle n \rangle/k)}\right), \end{aligned} \quad (18)$$

where $\hat{r}_1 \xrightarrow{\text{large } N} \frac{\langle n \rangle}{k} \left\{ 1 - \sqrt{1 - k(C_2 - 1 - 1/\langle n \rangle)} \right\} \equiv \frac{\langle n \rangle}{k} p$, and $\hat{r}_2 - \hat{r}_1 = \frac{\langle n \rangle}{N}$.

6. Concluding remarks

Equations (5) and (6) and the NBD are used in analyses of the data. We have found that several minimum χ^2 values are coincided with each other. Referring this fact, we have shown that Eq.(10) is derived from Eq.(5) as N is large. Moreover the generalized Glauber-Luchs formula is also derived from Eq.(17) as N is large.

We consider physical correspondences between QCD and multiparticle dynamics in Eq.(1). Using Eq.(4), $\exp(\lambda t) = 1 + r_2$, λ and t can be estimated by the data. They are shown in Fig. 2 and Table III. We obtain an interesting expressions for λ and t in

multiparticle dynamics. The different interval concerning t is probably attributed to the hadronization. More data at $\sqrt{s}=135$ GeV are necessary to confirm them. Finally it should be noticed that there is a different approach to stochastic equation in QCD in Ref. [22].

Acknowledgments: One of the authors (M. B.) is partially supported by the Grant-in Aid for Scientific Research from the Ministry of Education, Science, Sport and Culture (No. 06640383) and (No. 09440103).

References

- [1] M. Biyajima, Prog. Theor. Phys. **69**(1983)966; *ibid* **70**(1983)1468A; See references on the NBD cited therein.
- [2] M. Biyajima and N. Suzuki, Phys.Lett. **B143**(1984), 463.
- [3] B. Durand and I. Sarcevic, Phys. Lett. **B172**(1986),104; See also, D. Ellis, Proc. 11th SLAC Summer Institute, SLAC Report 267(1983).
- [4] R. C. Hwa and C. S.Lam, Phys. Lett. **B173** (1986),346.
- [5] M. Biyajima, T. Kawabe and N. Suzuki, Phys. Lett. **B189**(1988), 466; See also M.Biyajima, A.Bartl, T.Mizoguchi and N.Suzuki, *ibid* **B237** (1990) 563.
- [6] P. V. Chliapnikov and O. G. Tchikilev, Phys. Lett. **B242** (1990) 275;
- [7] N. Suzuki, M. Biyajima and G. Wilk, Phys. Lett. **B268** (1991) 447.
- [8] N. Suzuki, M. Biyajima and N. Nakajima, Phys. Rev. **D53** (1996) 3582 and **D54** (1996) 3653; See also *ibid* **D54** (1996) 4333, and J.E. Seger and J. Andrew Green, Phys. Lett **B351** (1995) 569.
- [9] P. V. Cliapnikov, O. G. Tchikilev and V. A. Uvarov, Phys. Lett. **B352** (1995) 461; O. G. Tchikilev, *ibid* **B382** (1996) 296 ; *ibid* **388**(1996),848.
- [10] T. Osada, N. Nakajima, M. Biyajima and N. Suzuki, DPSU-96-17 (unpublished).
- [11] J. Finkelstein, Phys. Rev. **D37** (1988) 2446.
- [12] V. Gupta and N. Sarma, Z. Phy. **C52** (1992) 53.
- [13] L. L. Chau and D. W. Huang, Phys. Rev. Lett. **70**(1993) 3380; See also L. L. Chau and D-W Huang, Phys. Lett.**B283**(1992) 1, D-W Hung, Mod. Phys. Lett. **A11** (1996) 2681.
- [14] HRS Collab., M. Derrick et al., Phys. Rev. **D34** (1986) 3304.

- [15] TASSO Collab., W. Braunschweig et al., Z. Phys. **C45** (1989) 193.
- [16] AMY Collab., H. W. Zheng et al., Phys. Rev. **D42** (1990) 737.
- [17] DELPHI Collab., P. Abreu et al., Z. Phys. **C52** (1991) 271.
- [18] OPAL Collab., P. D. Acton et al., Z. Phys. **C53** (1992) 539.
- [19] SLD Collab., K. Abe et al., Phys.lett. **B371** (1996) 149.
- [20] OPAL Collab., CERN-PPE/96-47.
- [21] H1 Collab., S. Aid et al., Z.Phys. **C72** (1997) 573.
- [22] P. Brogueira, J. Deas De Deus and I. P. Da Silva, in Proceedings of XXVII I-th Conf. on HEP (World-Scientific, 1996), 583.

Table captions

Table I:

The minimum values of χ^2 's for fitting of discrete distribution to the experimental data using the MNBD the NBD and the Laguerre-type distribution, Eq.(10), respectively. The data of $\langle n \rangle$ and C_2 are used. '*' denotes value of the minimum χ^2 's. The NBD is given by $P(n) = \Gamma(n+k)/\Gamma(n+1) \Gamma(k) (k/\langle n \rangle)^k (1+k/\langle n \rangle)^{-(n+k)}$.

Table II:

The same as Table I but for fitting of the KNO scaling function obtained from the MNBD, Eq. (7), the gamma distribution and the Laguerre-type distribution, Eq.(11), respectively. The gamma distribution is given by $\Psi(z) = k^k/\Gamma(k) e^{-kz} z^{k-1}$.

Table III:

Physical correspondences between λ and t in Eq.(1) in QCD and mutiparticle dynamics.

Figure Captions

Fig.1:

The result of fitting by NBD and MNBD to the data observed in e^+e^- collisions by DELPHI Collab.[14].

Fig.2:

$1+r_2 = \exp(\lambda t)$ vs $\sqrt{s}$. A function $\exp(\lambda \ln(\sqrt{s}/s_0))$ is chosen among various fittings by the MINUIT program: $\chi^2 = 28.9$. We can estimate λ and s_0 in multiparticle dynamics.

Exp.	[GeV]	n_{max}	[N]	MNBD	NBD	Eq.(10)
TASSO	14.	26	[9]	*9.34/10	25.0/11	25.0/11
TASSO	22.	28	[11]	*2.14/11	10.1/12	10.3/11
HRS	29.	28	[20]	*6.24/11	8.20/12	8.20/12
TASSO	34.8	36	[13]	*11.8/15	39.8/16	45.6/16
TASSO	43.6	38	[38]	7.42/16	7.40/17	*6.82/17
AMY	57.	40	[12]	*9.84/17	51.8/18	64.8/18
DELPHI	91.2	52	[10]	*14.2/22	185./23	278./23
OPAL	91.2	54	[9]	*4.03/24	47.5/25	76.2/25
SLD	91.2	50	[13]	*27.1/21	154./22	301./22
OPAL	133.	54	[7]	*4.57/21	39.6/22	57.2/22
H1 $1 < \eta^* < 5$	80-115	18	[18]	2.93/16	2.92/17	*2.32/17
	115-150	21	[21]	8.52/19	*7.01/20	7.80/20
	150-185	22	[22]	5.12/20	5.65/21	*4.75/21
	185-220	23	[23]	5.99/21	8.97/22	*5.53/22

Table I

Exp.	[GeV]	Eq.(7)	NBD	Eq.(11)
TASSO	14.	*10.9/11	19.0/12	*11.0/12
TASSO	22.	*3.53/12	15.3/13	*3.21/13
HRS	29.	15.0/12	37.4/13	*11.9/13
TASSO	34.8	*8.85/16	42.7/17	*9.23/17
TASSO	43.6	14.1/17	49.9/18	*12.3/18
AMY	57	*7.82/18	19.1/19	8.62/19
DELPHI	91.2	*17.1/23	21.0/24	141./24
OPAL	91.2	*4.46/25	*4.39/26	29.9/26
SLD	91.2	*35.3/22	200./23	85.0/23
OPAL	133	11.2/22	*9.63/23	44.9/23
H1 $1 < \eta^* < 5$	80-115	20.5/17	37.7/18	*18.9/18
	115-150	19.5/20	32.8/21	*18.6/21
	150-185	9.55/21	18.4/22	*9.04/22
	185-220	12.3/22	23.3/23	*11.7/23

Table II

Eq.(1)	P(n,t)	λ	λ_0	interval of t
QCD	gluon dis.	$\frac{N_c}{\varepsilon}$	$\frac{N_c^2-1}{2N_c\varepsilon}$	$\ln[\ln(Q^2/\Lambda^2)/\ln(Q_0^2/\Lambda^2)]$
multiparticle dynamics	hadron dis.	~ 0.36	0 (or $k = \lambda_0/\lambda$)	$\ln(\sqrt{s}/6.85)$

Table III




Fig. 1




Fig.2